

Promiscuity (continued)

relationships than in heterosexual marriage. In Holland, male homosexual relationships last an average of 1.5 years, and gay men have an average of eight partners a year outside of their supposedly “committed” relationships. (XyWrite M, et al. The contribution of steady and casual partnerships to the incidence of HIV infection among homosexual men in Amsterdam. AIDS. 2003; 17: 1029-38.)

Sexual Partners Often Unknown

The Kinsey Institute study revealed that 79% of the respondents said that over half of their sexual partners were strangers.

Violence

Gay authors, Island and Letellier, find much more violence exists in homosexual relationships than in male-female coupling. David Island and Patrick Letellier, *Men Who Beat The Men Who Love Them*, (New York, Haworth Press, 1991)

Suicide

Homosexual men are six times more likely to have attempted suicide than are heterosexual men. (Bell and Weinberg, *Homosexualities*, Table 21.12).

Drug and Alcohol Abuse

Studies find that between 25 and 33% of homosexual men and women are alcoholics while abuse of drugs, smoking and other high risk activities frequently

accompany homosexuality. (R. Kus, Alcoholics Anonymous and Gay American Men, Journal of Homosexuality, Volume 14, No.2 (1987), p.254).

Inability to Donate Blood

Males who have had sex with males, even once, are disqualified from giving blood in Hong Kong. Homosexual males cannot give blood because the risk is simply too high that the blood will contain an undetected infection. The health of the recipient of the blood is vital and is not risked with possibly infected blood. But what does this mean for the presence of disease in the blood of participants in homosexual sex?

Anal Intercourse

Anal intercourse, an extremely common practice among homosexual men, can seriously damage internal tissues in the anus and permanently weaken the anal sphincter, causing incontinence and other significant medical problems

Diseases from Anal Intercourse

A British study found that 60% of homosexual men engage in anal sex, frequently without condom and even if they know that they are HIV positive. (Mercer CH et al. Increasing prevalence of male homosexual partnerships and practices in Britain 1990-2000. AIDS. 2004; 18: 1453-8). As a result, a large number of diseases are associated with anal intercourse, many of which are rare or even unknown in the heterosexual population such as: anal cancer,

Chlamydia trachomatis, Cryptosporidium, Giardia lamblia, Herpes simplex virus, HIV, Human papilloma virus, Isospora belli, Microsporidia, Gonorrhoea, Syphilis, Hepatitis B and C and others. The risk of contracting HIV increases significantly among those engaging in anal intercourse.

Condom Use Flawed

Condoms do not protect fully protect users against HIV and fail as much as 15% of the time. Dr. Brian Clowes, *The Case Against Condoms*, U. S., Human Life International, 2006. Condoms, even when used 100% of the time, fail to give adequate levels of protection against sexually transmitted diseases such as Syphilis, Gonorrhoea, Chlamydia, Herpes, Genital Warts and others. (Sex, Condoms, and STDs: What We Now Know. Medical Institute for Sexual Health. 2002). Condoms used in the anus can create tears in anal tissue allowing fecal matter to spill into the bloodstream.

Sexually Transmitted Diseases

In a survey reported in an American Public Health Association publication, 78% of the gay respondents reported that they had been affected by a sexually transmitted disease at least one time. (Enrique T. Rueda, *The Homosexual Network*, Old Greenwich, Conn., The Devin Adair Company, 1982, p.53).

Longevity

In a study of 6,574 obituary notices for homosexual persons taken from 18 U. S. gay publications, the average age at death was 42, but it was 39 for those who died of AIDS. Only 9% of homosexual males reached age 65, but just 2% if they died of AIDS. This compares with 80% of married, heterosexual males reaching age 65. (Cameron, Playfair, and Wallum, *The Longevity of Homosexuals: Before and After the AIDS Epidemic*, Omega: Journal of Death and Dying, Vol. 29, #3, 1994, p. 252.)

Same-Sex Marriage

Allowing common law same-sex marriage in Scandinavia has diminished marriage as the ultimate standard there. This has greatly hastened the demise of the traditional family-based society. Births out of wedlock have risen to 50%. Marriage has declined, while unstable cohabitation has risen. The number of children who reach age 21 with both natural parents in the home has dramatically decreased. "Gay marriage is both an effect and a reinforcing cause of the separation of marriage and parenthood." Dr. Stanley Kurtz, *The End of Marriage in Scandinavia*, *The Weekly Standard* 02/02/2004, Volume 009, Issue 20

(www.weeklystandard.com/Content/Public/Articles/000/000/003/660zypwj.asp)

Sex Outside a Primary Relationship

Gay men have sex with someone other than their primary partner in 66% of relationships within the first year, rising to 90% of relationships after five years. (Joseph Harry, *Gay Couples*. (New York Praeger Press, 1984)

These figures are confirmed by gay authors McWhirter and Mattison. They studied 156 males couples and found over 95% of these participated in sex outside of the relationship. In addition, 89% masturbated. David McWhirter and Andrew Mattison, *The Male Couple: How Relationships Develop*, N. J., Prentice-Hall, 1984

With the use of three categories of sex, clearly gays in relationships practise sexual activities regularly. But there's no evidence of a biological imperative for this. People who have left homosexuality confirm that a person is NOT born gay.



For more information on homosexuality, health, other concerns or leaving homosexuality, please e-mail info@caringsfriends.hk or visit www.caringsfriends.hk

Facts from Books and Studies on Homosexuality

No Gay Gene

No replicated scientific studies in the world have found a genetic or biological origin for homosexuality.

Cause of Homosexuality

Ex-gay author, Leslie Lung states: "On the surface, homosexuality appears to be a sexual issue. It is NOT. Many issues are emotional ones, resulting from relationships gone wrong within the family." (*Freedom of Choice*, Lung, 2001)

Duration of Homosexual Relationships

Homosexual relationships are usually much shorter than heterosexual relationships. A Kinsey Institute study found that 78% of male homosexual "partnerships" lasted less than three years. Only 12% lasted five years or longer. (Bell and Weinberg, *Homosexualities: A Study of Diversity Among Men and Women*, (New York, Simon and Shuster, 1978, p.314).

Promiscuity

The same Kinsey study found that 75% of homosexual men admitted to having sex with more than 100 different males in their lifetime, while 28% of homosexual males surveyed claimed more than 1,000 lifetime sexual partners!

Far higher rates of promiscuity are observed even within 'committed' gay

Please turn over